

WEST**Freeform Search****Database:**

US Patents Full-Text Database
JPO Abstracts Database
EPO Abstracts Database
Derwent World Patents Index
IBM Technical Disclosure Bulletins

Term:

115 and 112

Display:

10

Documents in Display Format:

TI

Starting with Number

1

Generate: ☐ Hit List ☒ Hit Count ☐ Image

Search

Clear

Help

Logout

Interrupt

Main Menu

Show S Numbers

Edit S Numbers

Preferences

Search History**Today's Date:** 5/22/2000

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT,JPAB,EPAB,DWPI,TDBD	114 and 116	0	<u>L19</u>
USPT,JPAB,EPAB,DWPI,TDBD	115 and 114	0	<u>L18</u>
USPT,JPAB,EPAB,DWPI,TDBD	115 and 111	2	<u>L17</u>
USPT,JPAB,EPAB,DWPI,TDBD	115 and 112	68	<u>L16</u>
USPT,JPAB,EPAB,DWPI,TDBD	accase or ahas or (acetyl coa carboxylase) or (acetylcoa carboxylase) or (acetyl coacacboxylase) or acetylcoacacboxylase or acetohydroxyacid	878	<u>L15</u>
USPT,JPAB,EPAB,DWPI,TDBD	110 and 19	75	<u>L14</u>
USPT,JPAB,EPAB,DWPI,TDBD	110 and 18	12	<u>L13</u>
USPT,JPAB,EPAB,DWPI,TDBD	110 and 17	174	<u>L12</u>
USPT,JPAB,EPAB,DWPI,TDBD	110 and 16	101	<u>L11</u>
USPT,JPAB,EPAB,DWPI,TDBD	resistant or resistance	1812554	<u>L10</u>
USPT,JPAB,EPAB,DWPI,TDBD	11 and 15	361	<u>L9</u>
USPT,JPAB,EPAB,DWPI,TDBD	11 and 14	25	<u>L8</u>
USPT,JPAB,EPAB,DWPI,TDBD	11 and 13	489	<u>L7</u>
USPT,JPAB,EPAB,DWPI,TDBD	11 and 12	429	<u>L6</u>
USPT,JPAB,EPAB,DWPI,TDBD	fenoxaprop	588	<u>L5</u>
USPT,JPAB,EPAB,DWPI,TDBD	imazamox	35	<u>L4</u>
USPT,JPAB,EPAB,DWPI,TDBD	clethodim or sethoxydim	630	<u>L3</u>
USPT,JPAB,EPAB,DWPI,TDBD	fluazifop or quizalofop	620	<u>L2</u>
USPT,JPAB,EPAB,DWPI,TDBD	glyphosate OR (roundup or spator or muster or glifonox or glycel) OR (phosphonomethylglycine or ((phosphonomethyl or (phosphono methyl)) glycine))	3900	<u>L1</u>

WEST

Generate Collection

L16: Entry 6 of 68

File: USPT

Feb 22, 2000

DOCUMENT-IDENTIFIER: US 6028252 A
TITLE: Soybean variety 90B43

BSPR:

The present invention relates to a new and distinctive soybean variety, designated 90B43 which has been the result of years of careful breeding and selection. There are numerous steps in the development of any novel, desirable plant germplasm. Plant breeding begins with the analysis and definition of problems and weaknesses of the current germplasm, the establishment of program goals, and the definition of specific breeding objectives. The next step is selection of germplasm that possess the traits to meet the program goals. The goal is to combine in a single variety an improved combination of desirable traits from the parental germplasm. These important traits may include higher seed yield, resistance to diseases and insects, tolerance to drought and heat, and better agronomic qualities.

BSPR:

Backcross breeding has been used to transfer genes for simply inherited, highly heritable traits into a desirable homozygous variety or inbred line that is utilized as the recurrent parent. The source of the traits to be transferred is called the donor parent. After the initial cross, individuals possessing the desired traits of the donor parent are selected and repeatedly crossed (backcrossed) to the recurrent parent. The resulting plant is expected to have the attributes of the recurrent parent (e.g., variety) and the desirable traits transferred from the donor parent. This approach has been used extensively for breeding disease resistant varieties.

BSPR:

LDG=LODGING RESISTANCE. Lodging is rated on a scale of 1 to 9. A score of 9 indicates erect plants. A score of 5 indicates plants are leaning at a 45.degree. angle in relation to the ground and a score of 1 indicates plants are laying on the ground.

BSPR:

The oldest and most traditional method of analysis is the observation of phenotypic traits. The data is usually collected in field experiments over the life of the soybean plants to be examined. Phenotypic characteristics most often observed are for traits associated with seed yield, seed protein and oil content, lodging resistance, disease resistance, maturity, plant height, and shattering.

BSPR:

Soybean variety 90B43 is a purple flowered, soybean variety with gray pubescence and yellow hila. The variety exhibits outstanding yield potential. Soybean variety 90B43 exhibits multi-race Phytophthora tolerance as well as superior iron deficiency chlorosis resistance. Variety 90B43 possesses the Rps1C gene which confers resistance to races 1-3, 6-11, 13, 15, 17, 21, 23, 24, 26, 28-30, and 32. This Phytophthora resistance is unique for 90B43's very early maturity group. 90B43 further demonstrates excellent standability. The variety is particularly suited to the North Central, and Northern Plains, including the Red River Valley, Regions of the United States. There are few other varieties at this relative maturity and even fewer with the Phytophthora tolerance that this variety exhibits.

BSPR:

One commonly used selectable marker gene for plant transformation is the neomycin phosphotransferase II (nptII) gene, isolated from transposon Tn5, which when

placed under the control of plant regulatory signals confers resistance to kanamycin. Fraley et al., Proc. Natl. Acad. Sci. U.S.A., 80: 4803 (1983). Another commonly used selectable marker gene is the hygromycin phosphotransferase gene which confers resistance to the antibiotic hygromycin. Vanden Elzen et al., Plant Mol. Biol., 5: 299 (1985).

BSPR:

Additional selectable marker genes of bacterial origin that confer resistance to antibiotics include gentamycin acetyl transferase, streptomycin phosphotransferase, aminoglycoside-3'-adenyl transferase, the bleomycin resistance determinant. Hayford et al., Plant Physiol. 86: 1216 (1988), Jones et al., Mol. Gen. Genet., 210: 86 (1987), Svab et al., Plant Mol. Biol. 14: 197 (1990), Hille et al., Plant Mol. Biol. 7:171 (1986). Other selectable marker genes confer resistance to herbicides such as glyphosate, glufosinate or broxynil. Comai et al., Nature 317: 741-744 (1985), Gordon-Kamm et al., Plant Cell 2: 603-618 (1990) and Stalker et al., Science 242: 419-423 (1988).

BSPR:

Another class of marker genes for plant transformation require screening of presumptively transformed plant cells rather than direct genetic selection of transformed cells for resistance to a toxic substance such as an antibiotic. These genes are particularly useful to quantify or visualize the spatial pattern of expression of a gene in specific tissues and are frequently referred to as reporter genes because they can be fused to a gene or gene regulatory sequence for the investigation of gene expression. Commonly used genes for screening presumptively transformed cells include .beta.-glucuronidase (GUS), .beta.-galactosidase, luciferase and chloramphenicol acetyltransferase. Jefferson, R. A., Plant Mol. Biol. Rep. 5: 387 (1987), Teeri et al., EMBO J. 8: 343 (1989), Koncz et al., Proc. Natl. Acad. Sci. U.S.A. 84:131 (1987), De Block et al., EMBO J. 3: 1681 (1984).

BSPR:

(A) Plant disease resistance genes. Plant defenses are often activated by specific interaction between the product of a disease resistance gene (R) in the plant and the product of a corresponding avirulence (Avr) gene in the pathogen. A plant variety can be transformed with cloned resistance gene to engineer plants that are resistant to specific pathogen strains. See, for example Jones et al., Science 266: 789 (1994) (cloning of the tomato Cf-9 gene for resistance to Cladosporium fulvum); Martin et al., Science 262: 1432 (1993) (tomato Pto gene for resistance to Pseudomonas syringae pv. tomato encodes a protein kinase); Mindrinos et al., Cell 78: 1089 (1994) (Arabidopsis RSP2 gene for resistance to Pseudomonas syringae).

BSPR:

(B) A gene conferring resistance to a pest, such as soybean cyst nematode. See e.g. PCT Application WO96/30517; PCT Application WO93/19181.

BSPR:

(M) A hydrophobic moment peptide. See PCT application WO95/16776 (disclosure of peptide derivatives of Tachyplesin which inhibit fungal plant pathogens) and PCT application WO95/18855 (teaches synthetic antimicrobial peptides that confer disease resistance), the respective contents of which are hereby incorporated by reference.

BSPR:

(N) A membrane permease, a channel former or a channel blocker. For example, see the disclosure by Jaynes et al., Plant Sci. 89: 43 (1993), of heterologous expression of a cecropin-.beta., lytic peptide analog to render transgenic tobacco plants resistant to Pseudomonas solanacearum.

BSPR:

(O) A viral-invasive protein or a complex toxin derived therefrom. For example, the accumulation of viral coat proteins in transformed plant cells imparts resistance to viral infection and/or disease development effected by the virus from which the coat protein gene is derived, as well as by related viruses. See Beachy et al., Ann. Rev. Phytopathol. 28: 451 (1990). Coat protein-mediated resistance has been conferred upon transformed plants against alfalfa mosaic virus, cucumber mosaic virus, tobacco streak virus, potato virus X, potato virus Y, tobacco etch virus, tobacco rattle virus and tobacco mosaic virus. Id.

BSPR:

(S) A developmental-arrestive protein produced in nature by a plant. For example, Logemann et al., Bio/Technology 10: 305 (1992), have shown that transgenic plants expressing the barley ribosome-inactivating gene have an increased resistance to fungal disease.

BSPR:

(A) A herbicide that inhibits the growing point or meristem, such as an imidazalinone or a sulfonylurea. Exemplary genes in this category code for mutant ALS and AHAS enzyme as described, for example, by Lee et al., EMBO J. 7: 1241 (1988), and Miki et al., Theor. Appl. Genet. 80: 449 (1990), respectively.

BSPR:

(B) Glyphosate (resistance) imparted by mutant 5-enolpyruvyl-3-phosphikimate synthase (EPSP) and aroA genes, respectively) and other phosphono compounds such as glufosinate (phosphinothricin acetyl transferase, PAT) and Streptomyces hygroscopicus phosphinothricin-acetyl transferase, bar, genes), and pyridinoxy or phenoxy proprionic acids and cyclohexones (ACCase inhibitor-encoding genes). See, for example, U.S. Pat. No. 4,940,835 to Shah et al., which discloses the nucleotide sequence of a form of EPSP which can confer glyphosate resistance. A DNA molecule encoding a mutant aroA gene can be obtained under ATCC accession No. 39256, and the nucleotide sequence of the mutant gene is disclosed in U.S. Pat. No. 4,769,061 to Comai. European patent application No. 0 333 033 to Kumada et al. and U.S. Pat. No. 4,975,374 to Goodman et al. disclose nucleotide sequences of glutamine synthetase genes which confer resistance to herbicides such as L-phosphinothricin. The nucleotide sequence of a phosphinothricin-acetyl-transferase gene is provided in European application No. 0 242 246 to Leemans et al. De Greef et al., Bio/Technology 7: 61 (1989), describe the production of transgenic plants that express chimeric bar genes coding for phosphinothricin acetyl transferase activity. Exemplary of genes conferring resistance to phenoxy proprionic acids and cyclohexones, such as sethoxydim and haloxyfop, are the Accl-S1, Accl-S2 and Accl-S3 genes described by Marshall et al., Theor. Appl. Genet. 83: 435 (1992).

BSPU:

1. Genes That Confer Resistance To Pests or Disease And That Encode:

BSPU:

2. Genes That Confer Resistance To A Herbicide, For Example:

BSTL:

TABLE 1 _____ VARIETY DESCRIPTION INFORMATION
90B43 _____ A. Mature Seed Characteristics: Seed Coat Color: yellow Seed Coat Luster: dull Seed Size (grams per 100 seeds): 17 Hilum Color: yellow Cotyledon Color: yellow B. Leaf: Leaflet Shape: ovate Leaf Color: medium green C. Plant Characteristics: Flower Color: purple Pod Color: tan Plant Pubescence Color: Gray Plant Types: bushy Plant Habit: indeterminate Maturity Group: 04 D. Bacterial Diseases (S = susceptible R = resistant) E. Fungal Diseases (S = susceptible R = resistant) Phytophthora Rot (Phytophthora megasperma var. sojae): Race 1: R Race 2: R Race 3: R Race 4: S Race 7: R F. Viral Diseases (S = susceptible R = resistant) Bud Blight (Tobacco Ringspot Virus): S Yellow Mosaic (Bean Yellow Mosaic Virus): S Cowpea Mosaic (Cowpea Chlorotic Virus): S Pod Mottle (Bean Pod Mottle Virus): S Seed Mottle (Soybean Mosaic Virus): S G. Nematode Diseases (S = susceptible R = resistant) Soybean Cyst Nematode Race 3: S Iron Chlorosis: R Submitted Seed Content (% Protein) 35 Submitted Seed Content (% Oil) 19 _____ (PVP Certificate No.) is a Pioneer HiBred International, Inc. proprietary variety. Publications useful as references in interpreting Table 1 include: Caldwell, B. E. ed. 1973. "Soybeans: Improvement, Production, and Uses" Amer. Soc. Agron. Monograph No. 16; Buttery, B. R., and R. I. Buzzell 1968. "Peroxidase Activity in Seed of Soybean Varieties" Crop Sci. 8: 722-725; Hymowitz, T. 1973. "Electrophoretic analysis of SBTIA2 in the USDA Soybean Germplasm Collection" Crop Sci., 13: 420-421; Payne R. C., and L. F. Morris, 1976. "Differentiation of Soybean Varieties by Seedling Pigmentation Patterns" J. Seed. Technol. 1: 1-19. The disclosures of which are each incorporated by reference in their entirety

DEPR:

The results in table 2A compare Soybean variety 90B43 with another similarly

DEPR:

DEPR:

DEPR:

The results in table 2D compare Soybean variety 90B43 with another similarly adapted Pioneer brand soybean variety, 9042. The results show that variety 90B43 is significantly higher yielding than variety 9042. Variety 90B43 also demonstrates significantly superior resistance to lodging than variety 9042. While not specifically shown in the table, Variety 9007 possesses the Rps1A gene which confers different races of Phytophthora resistance (races 1, 2, 10, 11, 13-18, 24, 26, 27 31, and 32) than the Rps1C gene possessed by variety 90B43 (races 1-3, 6-11, 13, 15, 17, 21, 23, 24, 26, 28-30, and 32).

22428 SEARCH REQUEST FORM

Examiner # (Mandatory): 69462 Requester's Full Name: Mark C. Gird
 Art Unit 1616 Location (Bldg/Room#): CM1-2D11 Phone (circle 305 306 308) 4550
 Serial Number: 09/264,775 Results Format Preferred (circle): PAPER DISK E-MAIL
 Title of Invention _____
 Inventors (please provide full names): attached
 Earliest Priority Date: _____

Keywords (include any known synonyms registry numbers, explanation of initialisms):

AHAS = acetoxyhydroxy acid synthase 9027-45-6
 ACCase = acetyl Co-A carboxylase 9023-93-2
glyphosate tolerance

RECEIVED
 NOV 27 1991
 CH/CHEN, ONI (STIC)
 claims & biblio. data
 attached

Search Topic:

Please write detailed statement of the search topic, and the concept of the invention. Describe as specifically as possible the subject matter to be searched. Define any terms that may have a special meaning. Give examples of relevant citations, authors, etc., if known. You may include a copy of the abstract and the broadcast or most relevant claim(s).

Herbicidal composition comprising:

- 1) glyphosate
- 2) a 2nd herbicide which inhibits either ACCase or AHAS:
 i.e.: flazasitop, quizalofop, clethodim, sethoxydim,
imazamox, fenoxaprop

Methods of use in glyphosate tolerant crops (cl. 9)

STAFF USE ONLY

Searcher: 22428
 Searcher Phone #: 44488
 Searcher Location: _____
 Date Picked Up: 12/11
 Date Completed: 12/13
 Clerical Prep Time: 10
 Terminal Time: 25/45
 Number of Databases: 4

Type of Search
 _____ N.A. Sequence
 _____ A.A. Sequence
☒ Structure (#)
 _____ Bibliographic
 _____ Litigation1
 _____ Fulltext
 _____ Procurement
 _____ Other

Vendors (include cost where applicable)
☒ STN
 _____ Questel/Orbit
 _____ Lexis/Nexis
 _____ WWW/Internet
 _____ In-house sequence systems (list)
 _____ Dialog
 _____ Dr. Link
 _____ Westlaw
 _____ Other (specify)

=> d his

(FILE 'HOME' ENTERED AT 09:12:56 ON 13 DEC 1999)

SET COST OFF
SET AUHELP OFF

FILE 'REGISTRY' ENTERED AT 09:13:31 ON 13 DEC 1999

L1 2 S 9027-45-6 OR 9023-93-2
E GLYPHOSATE/CN

L2 1 S E3
E C3H8NO5P/MF
E FLUAZIFOP/CN

L3 1 S E3
E C15H12F3NO4/MF

L4 13 S E3 AND NC5/ES AND 46.150.18/RID AND 2/NR

L5 2 S L4 AND (2R OR S)

L6 3 S L3,L5
E QUIZALOFOP/CN

L7 1 S E3
E C17H13CLN2O4/MF

L8 9 S E3 AND NC2NC2-C6/ES AND 46.150.18/RID AND 3/NR

L9 7 S L8 NOT (ESTER OR QUINOXALINECARBOXYLIC)

L10 4 S L9 NOT 6 CHLORO

L11 1 S L10 AND IDS/CI

L12 3 S L9 NOT L10

L13 4 S L7,L11,L12
E CLETHODIM/CN

L14 1 S E3
E C17H26CLNO3S/MF

L15 12 S E3 AND C6/ES AND 1/NR

L16 4 S L15 NOT 46.150.18/RID

L17 2 S L16 NOT 3 CYCLOHEX?

L18 2 S L14,L17
E SETHOXYDIM/CN

L19 1 S E3
E C17H20NO3S/MF
E C17H29NO3S/MF

L20 8 S E3 AND C6/ES NOT 46.150.18/RID

L21 3 S L20 NOT 3 CYCLOHEX?

L22 2 S L21 NOT SC5/ES
E IMAZAMOX/CN

L23 1 S E3
E C15H19N3O4/MF

L24 13 S E3 AND NCNC2/ES AND NC5/ES AND 2/NR

L25 3 S L24 AND METHOXYMETHYL

L26 2 S L25 NOT 6

L27 2 S L23,L26
E FENOXAPROP/CN

L28 1 S E3

L29 3 S 1113776-21-9 OR 113158-40-0 OR 95617-09-7 OR 73519-45-6

L30 2 S L29 NOT 5 CHLORO

L31 2 S L28,L30

L32 1 S 113776-21-9

L33 3 S L31,L32

L34 14 S L6,L12,L18,L19,L27,L33
SAV L34 CLARDY264/A

FILE 'HCAOLD' ENTERED AT 09:34:49 ON 13 DEC 1999

L35 0 S L34

FILE 'HCAPLUS' ENTERED AT 09:35:00 ON 13 DEC 1999

L36 1242 S L34

L37 1449 S FLUAZIFOP OR QUIAZLOFOP OR CLETHODIM OR SETHOXYDIM OR IMAZAMO

L38 1708 S L36,L37

L39 2470 S L1

L40 601 S (ACETOLACTATE OR ACETOLACTIC OR ACETO() (LACTATE OR LACTATE))

L41 275 S (ACETOHYDROXY OR ACETO HYDROXY) () ACID () (SYNTHETASE OR SYN
L42 178 S ACETOHYDROXYACID () (SYNTHETASE OR SYNTHASE)
L43 2039 S ACETYL () (COA OR COENZYME A) () CARBOXYLASE
L44 3131 S L39-L43
L45 4035 S L2 OR GLYPHOSATE
L46 16 S L39 AND L45
L47 16 S L44 AND L46
L48 659 S L39 AND (SYNERG? OR MIX? OR COMBIN? OR COMPOSITION)
L49 659 S L48 AND L44
L50 3 S L48 AND L45
L51 3 S L49 AND L50
E FLINT J/AU
L52 24 S E3,E7,E14,E15
E PROBST N/AU
L53 7 S E3,E7
E GUBBIGA N/AU
L54 5 S E4
L55 1 S L52-L54 AND L38
L56 0 S L52-L54 AND L44
L57 3 S L52-L54 AND L45
L58 3 S L55,L57
L59 2 S L51 NOT MRNA/TI
L60 2 S L45 AND NONGLYPHOSATE
L61 1439 S L45 AND (MIX? OR SYNERG? OR COMPOSITION OR COMBIN? OR FORMUL?
L62 6 S L61 AND L44
L63 6 S L62 NOT L58,L60
L64 5 S L63 NOT MRNA/TI
L65 10 S L58,L60,L64

FILE 'HCAPLUS' ENTERED AT 09:48:23 ON 13 DEC 1999
SEL RN L58

FILE 'REGISTRY' ENTERED AT 09:49:12 ON 13 DEC 1999

L66 16 S E1-E16
L67 1 S 81591-80-2
L68 2 S L2,L67
SEL RN
L69 577 S E17-E18/CRN
SEL RN L34
L70 374 S E19-E32/CRN
L71 0 S L68 AND L70
L72 8 S L69 AND L70

FILE 'HCAOLD' ENTERED AT 09:51:44 ON 13 DEC 1999

L73 0 S L72

FILE 'HCAPLUS' ENTERED AT 09:51:46 ON 13 DEC 1999

L74 3 S L72
L75 5 S L74,L58

FILE 'USPATFULL' ENTERED AT 09:52:09 ON 13 DEC 1999

L76 0 S L72

=> fil hcaplus

FILE 'HCAPLUS' ENTERED AT 09:52:35 ON 13 DEC 1999

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 1999 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE COVERS 1967 - 13 Dec 1999 VOL 131 ISS 25

FILE LAST UPDATED: 12 Dec 1999 (19991212/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

This file supports REGISTRY for direct browsing and searching of all substance data from the REGISTRY file. Enter HELP FIRST for more information.

=> d all tot 175 hitstr

L75 ANSWER 1 OF 5 HCAPLUS COPYRIGHT 1999 ACS
 AN 1999:594855 HCAPLUS
 DN 131:195769
 TI Mixtures for weed control in **glyphosate**-tolerant soybean
 IN **Flint, Jerry L.; Probst, Norman J.; Gubbiga, Nagabhushana G.**
 PA Monsanto Company, USA
 SO PCT Int. Appl., 38 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 IC ICM A01N057-20
 ICS A01N057-20; A01N043-76; A01N043-60; A01N043-50; A01N043-40;
 A01N035-10

CC 5-3 (Agrochemical Bioregulators)

FAN 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9945781	A1	19990916	WO 1999-US5089	19990309
W:				
AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW:				
GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				

PRAI US 1998-PV77241 19980309

AB The present invention is directed to tank mixts. and premixts. of a **glyphosate** herbicide and a second herbicide to which a first species is susceptible and a second species is resistant. Such tank mixts. and premixts. allow control of **glyphosate**-susceptible weeds and **glyphosate**-tolerant volunteer individuals of the first species in a crop of **glyphosate**-tolerant second species with a single application of herbicide. Particularly, the invention relates to the control of volunteer **glyphosate**-tolerant corn in a crop of **glyphosate**-tolerant soybean.

ST weed control **glyphosate** tolerant soybean

IT Soybean (Glycine max)

(**glyphosate**-tolerant; weed control in)

IT Weed control (herbicide)

(mixts. for weed control in **glyphosate**-tolerant soybean)

IT Corn

(volunteer; control in **glyphosate**-tolerant soybean)

IT 1071-83-6D, **Glyphosate**, mixts. contg.

242132-22-5, **Glyphosate-fluazifop** mixt.

242132-23-6, **Glyphosate-quizalofop** mixt.

242132-24-7 242132-25-8 242132-26-9

242132-27-0 242132-28-1 242143-59-5, Fusilade DX-Touchdown mixt.

RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses)
 (weed control in **glyphosate**-tolerant soybean by)


IT 1071-83-6D, **Glyphosate**, mixts. contg.

242132-22-5, Glyphosate-fluazifop mixt.
 242132-23-6, Glyphosate-quizalofop mixt.
 242132-24-7 242132-25-8 242132-26-9
 242132-27-0

RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses)
 (weed control in **glyphosate**-tolerant soybean by)

RN 1071-83-6 HCAPLUS

CN Glycine, N-(phosphonomethyl)- (7CI, 8CI, 9CI) (CA INDEX NAME)




RN 242132-22-5 HCAPLUS

CN Glycine, N-(phosphonomethyl)-, mixt. with 2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoic acid (9CI) (CA INDEX NAME)

CM 1

CRN 69335-91-7


CMF C15 H12 F3 N O4



CM 2

CRN 1071-83-6

CMF C3 H8 N O5 P




RN 242132-23-6 HCAPLUS

CN Glycine, N-(phosphonomethyl)-, mixt. with 2-[4-[(6-chloro-2-quinoxalinyloxy]phenoxy]propanoic acid (9CI) (CA INDEX NAME)

CM 1

CRN 76578-12-6


CMF C17 H13 Cl N2 O4



CM 2

CRN 1071-83-6

CMF C3 H8 N O5 P



RN 242132-24-7 HCAPLUS


CN Glycine, N-(phosphonomethyl)-, mixt. with 2-[1-[[[3-chloro-2-propenyl)oxy]imino]propyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one (9CI) (CA INDEX NAME)

CM 1

CRN 99129-21-2

CMF C17 H26 Cl N O3 S


CDES *



CM 2

CRN 1071-83-6

CMF C3 H8 N O5 P




RN 242132-25-8 HCAPLUS

CN Glycine, N-(phosphonomethyl)-, mixt. with 2-[1-(ethoxyimino)butyl]-5-[2-(ethylthio)propyl]-3-hydroxy-2-cyclohexen-1-one (9CI) (CA INDEX NAME)

CM 1

CRN 74051-80-2


CMF C17 H29 N O3 S



CM 2

CRN 1071-83-6

CMF C3 H8 N O5 P




RN 242132-26-9 HCAPLUS

CN Glycine, N-(phosphonomethyl)-, mixt. with 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic acid (9CI) (CA INDEX NAME)

CM 1

CRN 114311-32-9

CMF C15 H19 N3 O4



CM 2

CRN 1071-83-6

CMF C3 H8 N O5 P




RN 242132-27-0 HCAPLUS

CN Glycine, N-(phosphonomethyl)-, mixt. with 2-[4-[(6-chloro-2-benzoxazolyl)oxy]phenoxy]propanoic acid (9CI) (CA INDEX NAME)

CM 1

CRN 95617-09-7


CMF C16 H12 Cl N O5



CM 2

CRN 1071-83-6

CMF C3 H8 N O5 P



L75 ANSWER 2 OF 5 HCAPLUS COPYRIGHT 1999 ACS

AN 1999:561282 HCAPLUS

DN 131:224816


TI Purple nutsedge (Cyperus rotundus) and sicklepod (Senna obtusifolia) response to glyphosate mixtures with ALS-inhibiting herbicides

AU Rao, A. Subrahmanyeswara; Reddy, Krishna N.
 CS Food and Agriculture Organization Fellow and Plant Physiologist, Southern,
 Agricultural Research Service, Stoneville, MS, 38776, USA
 SO Weed Technol. (1999), 13(2), 361-366
 CODEN: WETEE9; ISSN: 0890-037X
 PB Weed Science Society of America
 DT Journal
 LA English
 CC 5-3 (Agrochemical Bioregulators)
 AB Greenhouse studies were conducted to evaluate potential interactions among
 glyphosate mixts. with five acetolactate synthase (ALS)-inhibiting
 herbicides (chlorimuron, imazamox, imazaquin, MON 12,000, or pyriithiobac)
 for the control of purple nutsedge and sicklepod at two growth stages.
 Herbicides were tested alone at 0.5.times. and 1.times. rates (1.times.
 being suggested use rate for these herbicides) and in combination with
 glyphosate at 560 (0.5.times.) and 1,120 (1.times.) g/ha on 3-wk-old
 plants and at 1,120 g/ha on 6-wk-old plants. Glyphosate alone, at 1,120
 g/ha, gave complete control of purple nutsedge and at least 78% control of
 sicklepod regardless of growth stage. In 3-wk-old purple nutsedge plants,
 3 of the 20 herbicide combinations were antagonistic and 17 combinations
 were additive, whereas all 5 combinations were additive in 6-wk-old
 plants. In sicklepod, 8 combinations were antagonistic and 12
 combinations were additive in 3-wk-old plants, and all 5 combinations were
 antagonistic in 6-wk-old plants. In 3-wk-old plants, the glyphosate
 (0.5.times.) plus imazaquin (0.5.times.) combination resulted in highest
 antagonism in purple nutsedge control (79%), and the combination of
 glyphosate (0.5.times.) plus imazamox (0.5.times.) resulted in highest
 antagonism in sicklepod control (54%). Mixing chlorimuron, imazamox,
 imazaquin, MON 12,000, or pyriithiobac with glyphosate does not increase
 glyphosate efficacy on purple nutsedge or sicklepod.
 ST Cyperus Senna glyphosate herbicide mixt
 IT Cyperus rotundus
 Senna (Cassia tora)
 Weed control (herbicidal)
 (Cyperus rotundus and Senna obtusifolia response to glyphosate mixts.
 with ALS-inhibiting herbicides)
 IT 1071-83-6, Glyphosate 142275-97-6, Glyphosate-chlorimuron mixt.
 142275-98-7, Glyphosate-imazaquin mixt. **242132-26-9**,
 Glyphosate-imazamox mixt. 244061-34-5, Glyphosate-MON 12000 mixt.
 244061-36-7, Glyphosate-pyriithiobac mixt.
 RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses)
 (Cyperus rotundus and Senna obtusifolia response to glyphosate mixts.
 with ALS-inhibiting herbicides)
 IT **242132-26-9**, Glyphosate-imazamox mixt.
 RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses)
 (Cyperus rotundus and Senna obtusifolia response to glyphosate mixts.
 with ALS-inhibiting herbicides)
 RN 242132-26-9 HCAPLUS
 CN Glycine, N-(phosphonomethyl)-, mixt. with 2-[4,5-dihydro-4-methyl-4-(1-
 methylethyl)-5-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-
 pyridinecarboxylic acid (9CI) (CA INDEX NAME)

CM 1

CRN 114311-32-9


CMF C15 H19 N3 O4



CM 2

CRN 1071-83-6

CMF C3 H8 N O5 P



L75 ANSWER 3 OF 5 HCAPLUS COPYRIGHT 1999 ACS

AN 1999:215560 HCAPLUS

DN 130:233650

TI Synergistic herbicidal compositions

IN De Carvalho Castro, Kelly Neoob; Mendonca, Wilson; Malefyt, Timothy; Salzman, Frederick P.; Watkins, Robert M.

PA American Cyanamid Company, USA

SO PCT Int. Appl., 48 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A01N057-20

ICS A01N025-02; A01N057-20; A01N043-50

CC 5-3 (Agrochemical Bioregulators)

FAN. CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9913723	A1	19990325	WO 1998-US18981	19980914
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	AU 9894791	A1	19990405	AU 1998-94791	19980914
	JP 11246316	A2	19990914	JP 1998-276402	19980914
PRAI	US 1997-936186		19970917		
	WO 1998-US18981		19980914		
AB	Synergistic control of Ipomoea, Cyperus, Sida and Euphorbia, etc., comprises applying a combination of glyphosate and an imidazolinone deriv., such as imazethapyr, imazaquin, imazapic, imazamox and imazapyr. Further provided are synergistic herbicidal compns. comprising glyphosate and an imidazolinone compd., and specifically concd. aq. herbicidal compns. of imidazolinyl acid salts and glyphosate salts.				
ST	synergism herbicide compn glyphosate imidazolinone deriv				
IT	Synergistic herbicides (compns. contg. glyphosate and imidazolinone deriv.)				
IT	Cyperus Euphorbia Ipomoea Sida (control by synergistic herbicidal compns. contg. glyphosate and imidazolinone deriv.)				
IT	1071-83-6D, Glyphosate, mixt. with imidazolinone derivs. 221298-59-5, Roundup-imazethapyr mixt.. 221298-60-8, Roundup-imazapic mixt. 221298-61-9, Roundup-imazamox mixt. 221298-63-1 221298-65-3 221298-67-5 221321-46-6 221321-51-3 RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses) (synergistic herbicidal compn.)				
IT	221298-61-9, Roundup-imazamox mixt. 221298-65-3 RL: AGR (Agricultural use); BIOL (Biological study); USES (Uses) (synergistic herbicidal compn.)				


RN 221298-61-9 HCAPLUS

CN Glycine, N-(phosphonomethyl)-, compd. with 2-propanamine (1:1), mixt. with 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic acid (9CI) (CA INDEX NAME)

CM 1

CRN 114311-32-9

CMF C15 H19 N3 O4



CM 2


CRN 38641-94-0

CMF C3 H9 N . C3 H8 N O5 P

CM 3

CRN 1071-83-6


CMF C3 H8 N O5 P



CM 4

CRN 75-31-0

CMF C3 H9 N



RN 221298-65-3 HCAPLUS


CN Glycine, N-(phosphonomethyl)-, compd. with 2-propanamine (1:1), mixt. with 2-[(4R)-4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic acid (9CI) (CA INDEX NAME)

CM 1

CRN 221298-64-2

CMF C15 H19 N3 O4

Absolute stereochemistry.



CM 2


CRN 38641-94-0

CMF C3 H9 N . C3 H8 N O5 P

CM 3

CRN 1071-83-6


CMF C3 H8 N O5 P



CM 4

CRN 75-31-0

CMF C3 H9 N



L75 ANSWER 4 OF 5 HCAPLUS COPYRIGHT 1999 ACS

AN 1990:32084 HCAPLUS

DN 112:32084

TI Antagonism of **glyphosate** toxicity to johnsongrass (*Sorghum halepense*) by 2,4-D and dicambaAU **Flint, Jerry L.**; Barrett, Michael

CS Dep. Agron., Univ. Kentucky, Lexington, KY, 40545-009, USA

SO Weed Sci. (1989), 37(5), 700-5

CODEN: WEESA6; ISSN: 0043-1745

DT Journal

LA English

CC 5-3 (Agrochemical Bioregulators)

AB Greenhouse studies were conducted to det. the basis for reduced johnson grass control when **glyphosate** was applied in mixts. with 2,4-D or dicamba. **Glyphosate** was applied to johnson grass at 0.28, 0.56, 0.84, and 1.12 kg/ha alone and in combination with 2,4-D or dicamba at 0.14, 0.28, 0.14, or 0.56 kg/ha. Johnson grass shoot and root fresh wts. measured 4 wk after treatment were higher when **glyphosate** was applied with 2,4-D (0.28 kg/ha **glyphosate**) or dicamba (0.28 kg/ha or 0.56 kg/ha **glyphosate**) compared to **glyphosate** applied alone at these rates. The antagonism of johnson grass control was not obsd. with combinations of some of the higher **glyphosate** rates with 2,4-D (0.56 or 0.84 kg/ha **glyphosate**) or dicamba (0.84 or 1.12 kg/ha **glyphosate**). The redn. of **glyphosate** activity on johnson grass occurred when any of four forms of 2,4-D or two forms of dicamba were added to the **glyphosate** spray mixt. **Glyphosate** uptake into johnson grass leaves and subsequent

translocation to the roots was reduced by the presence of 2,4-D or dicamba. The reduced **glyphosate** uptake and translocation could account for the decreased toxicity of **glyphosate** to johnson grass when applied with 2,4-D or dicamba.

ST johnson grass control glyphosphate dichlorophenoxyacetate dicamba
 IT Johnson grass
 (control of, by glyphosphate, 2,4-D and dicamba antagonism of)
 IT Weed control
 (of johnson grass, by glyphosphate, 2,4-D and dicamba antagonism of)
 IT 94-75-7, 2,4-D, biological studies 1918-00-9, Dicamba
 RL: BIOL (Biological study)
 (johnson grass control by glyphosphate antagonism by)
 IT 1071-83-6
 RL: BIOL (Biological study)
 (johnson grass control by, 2,4-D and dicamba antagonism of)
 IT 1071-83-6
 RL: BIOL (Biological study)
 (johnson grass control by, 2,4-D and dicamba antagonism of)
 RN 1071-83-6 HCAPLUS
 CN Glycine, N-(phosphonomethyl)- (7CI, 8CI, 9CI) (CA INDEX NAME)

HO₂C-CH₂-NH-CH₂-PO₃H₂

L75 ANSWER 5 OF 5 HCAPLUS COPYRIGHT 1999 ACS

AN 1989:149765 HCAPLUS

DN 110:149765

TI Effects of **glyphosate** combinations with 2,4-D or Dicamba on field bindweed (*Convolvulus arvensis*)

AU Flint, Jerry L.; Barrett, Michael

CS Dep. Agron., Univ. Kentucky, Lexington, KY, 40546-0091, USA

SO Weed Sci. (1989), 37(1), 12-18

CODEN: WEESA6; ISSN: 0043-1745

DT Journal

LA English

CC 5-3 (Agrochemical Bioregulators)

AB Applications of isopropylamine **glyphosate** at 0.28, 0.56, 0.84, and 1.12 kg active ingredient/ha in combination with the dimethylamine salts of 2,4-D or dicamba at 0.14, 0.28, 0.41, and 0.56 kg active ingredient/ha produced additive or synergistic field bindweed control compared to the herbicides applied alone. Leaf and root growth was inhibited more from herbicide combinations than would be predicted from the effects of the chems. applied alone at the same rate. The uptake of ¹⁴C from **glyphosate** into the treated leaf and its accumulation in roots increased when 2,4-D or dicamba was combined with the 0.28 kg/ha rate of [¹⁴C]**glyphosate**. The combination of 2,4-D or dicamba with a higher (0.84 kg/ha) [¹⁴C]**glyphosate** rate did not change total absorption of ¹⁴C from **glyphosate**. However, compared to 0.84 kg/ha of [¹⁴C]**glyphosate** applied alone, less ¹⁴C accumulated above the treated leaf and more accumulated in the roots when 2,4-D was added to the **glyphosate**. The combination of **glyphosate** with 2,4-D or dicamba generally resulted in both increased uptake of ¹⁴C from 2,4-D or dicamba and greater accumulation in the roots. The additive or synergistic field bindweed control obsd. from mixts. of **glyphosate** with 2,4-D or dicamba appeared to be due to greater accumulation of the herbicides in the roots.

ST **glyphosate** dichlorophenoxyacetate dicamba field bindweed control

IT *Convolvulus arvensis*

 (control of, by **glyphosate** combination with 2,4-D or dicamba)

IT Weed control

 (of field bindweed, with **glyphosate** combinations with 2,4-D or dicamba)

IT Biological transport

 (absorption, of **glyphosate** combinations with 2,4-D or

IT 2008-39-1 2300-66-5, Dicamba dimethylamine salt 38641-94-0
75547-81-8 75553-94-5
RL: BIOL (Biological study)
(field bindweed control by)